ANESTHETIC EMERGENCIES & ACCIDENTS

LYON LEE, DVM PHD DACVA

Human Errors

- Miscalculation of the drug dose, e.g. getting one decimal wrong can mean ten times of overdose that may induce severe toxic effect
- Mislabeling of the syringe, misfiling to a wrong vaporizer etc. may constitute severe hazard
- Equipment failure or misused devices, most notably delivery of hypoxic mixture to the patient
- Exercise precautions to avoid human induced medical mishaps

Bradycardia

- As a general rule for heart rate < 60 beats per min in dogs x

  Parasympathetic stimulation (distend bladder, manipulation of the viscera, oculocardiac reflex etc.)

Problem

- Reduced cardiac output
- Arrhythmias (danger to progress to worse arrhythmias, e.g. complete cardiac arrest, if not corrected timely)

Treatment

- Identify cause and address it
- Dog and cat:
  - Atropine – 0.02 ~ 0.04 mg/kg IV
  - Glycopyrrolate – 0.005 ~ 0.01 mg/kg IV
  - If no response to above treatment, then consider giving isoproterenol 0.5 mcg/kg or epinephr0 0 12 126[J 7 0 T0.t6 7 0 T0M10(I)13(V)]TJ 5.8V-2(bnol)-2usT0.t63.497 0 Td ( )T

Brachycephalic Problems

Anatomical handicaps

- Stenotic nares
- Elongated soft palate
- Everted laryngeal ventricles
- Hypoplastic trachea

produce maximal flow. Release the pressure completely during relaxation to allow cardiac filling.

- Blood flow generated by cardiac compression will temporarily sustain cerebral and myocardial viability only if oxygenation is adequate. However, cerebral blood flow is less than one-fifth normal and coronary flow even less during external compression.
- Complications, which may arise from external compression, include sternal and rib fractures, blunt traumatic damage to intrathoracic and abdominal viscera and pneumothorax.

Mechanisms of blood flow during external chest compression

- Controversial.
- Two theories predominate. They probably both operate.
  - *Cardiac pump theory*. External chest compression squeezes the heart between t

- Epinephrine
  - Benefits are mainly due to its alpha-adrenergic actions.
  - It increases arterial wall tone and total peripheral resistance. This allows better intrathoracic arterial flow by reducing the tendency of vessels to collapse from the pressure induced by chest compressions.
  - Epinephrine also increases diastolic pressure which is important for myocardial perfusion, and renders the fibrillating heart more apt to defibrillate
  - It also diverts blood flow away from non-vital, towards vital tissues.
  - Dose of epinephrine is between 0.02 mg/kg to 0.2 mg/kg. Repeat this dose each 3 to 5 minutes if needed
- Sodium bicarbonate
  - Administered when a metabolic acidosis has developed or is strongly suspected (indicated by the patient's history) or if resuscitation has proceeded for longer than 20 minutes.
  - The dose is approximately 0.5-1 mEq/kg for each 10 minutes of arrest.
  - Venous blood gases are most indicative of bicarbonate needs.
  - Bicarbonate therapy is currently controversial, with concern that overzealous use may induce hyperosmolality, hypernatremia, paradoxical intracellular acidosis, and iatrogenic alkalosis. (see Acid Base Physiology and Anesthesia lecture) Alkalosis lowers the serum potassium levels and shifts the oxy-hemoglobin dissociation

- Inotropic support (e.g. dopamine at 5-10 mcg/kg min intravenous infusion) to maintain blood pressure
- Aim is to optimize cerebral perfusion and oxygen delivery (maintain normotension, moderate hypoventilation to PCO_2 of 30 ~ 35 mm Hg, maintain normal intracranial pressure with diuretics and corticosteroids, oxygen therapy) and control the cerebral metabolic demand for oxygen (suppress seizures). Barbiturates and hypothermia have not been shown to be useful under these circumstances. Neuroprotective drugs such as free radical scavengers (DMSO, desferoxamine) and calcium channel blockers are, at this time, of experimental value only and result in poor to variable responses.
- Close monitoring at least for further 24 hours is strongly recommended after successful CPR.