PERIOPERATIVE PAIN MANAGEMENT

Lyon Lee DVM PhD DACVA

Introduction

- **Pain** is defined as an unpleasant sensory and emotional experiences associated with actual or potential tissue damage or described in terms of such damage (Merskey 1979).
- Pain is a universal experience for every living being, and we all know what it feels like, but it is still very difficult to clearly define or quantify.
- Pain management in animals at perioperative and postoperative periods presents a significant challenge not only to veterinarians but also to the owners of the animals.
- Humans can express and describe the sensations they experience, and these descriptors are well accepted. However, in animals the difficulty of obtaining feedback through communication complicates the management of pain, as one can only attempt to help alleviate the pain when it is expressed.
- It is, therefore very important that one is familiar with the signs associated with pain and interpretation of these behavioral changes for the benefit of clinical pain management in animals.
- The ability to feel pain has clear survival advantages;
  - limits movement of the damaged part so that it promotes healing
  - the individuals learn to avert similar noxious stimuli in the future
- However, ongoing pain has no benefit and offers disadvantages;
  - Animals feeling pain postoperatively get distressed and the stress response interferes with normal resumption of activity
  - Delay in wound healing has been demonstrated in animals in more painful condition
  - Decreased food intake impairs nutrition and speed of recovery from the operation
  - Impaired respiration, especially in animals undergoing thoracic surgery, contributes to development of hypoxia, pulmonary atelectasis, pneumothorax, and retention of mucus and sputum etc.
  - May self-mutilate
  - Causes central sensitization which is more resistant to conventional analgesic therapy
- The reasons for relative reluctance in treating pain have been;
  - The clinician is not absolutely sure whether the animals are in pain
  - Belief of over-rated concerns regarding the pain and subsequent over-prescription of analgesics is dangerous due to its adverse side effects
  - Many effective analgesics are controlled substances and require a tight inventory log
  - Belief of analgesics should be withheld so that protective function of pain is maintained.
  - Insufficient knowledge about the analgesics, especially regarding opioids.
- There is no doubt that a skiob2ooa. l2t p0 1 Tf 16 mcri(s)-1()4(s)-1-6(bout)cmpe(no viht)-2(3m [(M)-ou) role in the management of suffering in animals.
- In addition, sound knowledge in physiology of pain and pharmacology of pain control substances (analgesics) will be valuable to maximize the therapeutic outcome in animals.

- Acute postoperative pain accompanies signs of guarding and aversion of the site, particularly

- Local analgesics
- General anesthetic. (NB, unconsciousness does not mean that all response to pain is removed; some general anesthetics are poor analgesics).
- Opioids
- NSAIDs
- Alpha 2 adrenoceptor agonists
- NMDA antagonists
- Adjunctive methods (unconventional analgesics)
- Choice of suitable one depends on type of pain, ease of use (how given) speed of effect, duration of

Opioids

- Opium (the plant) has been used for its analgesic and euphoric effects over thousands of years for men.
- All opioids (referring all endogenous and exogenous substances that possess morphine-like properties) are chemically related and subject to regulatory controls, governed by the Controlled Substance Act 1970
- In man and dog these are the mainstay of perioperative analgesia, and can be used in all three phases; pre/intra/post-operative periods.
- The potential excitement in cat and horse limit the use of the very potent drugs.
- The choice of opioid suitable for each phases is based on the following points.
- Premedication
    - This needs a long acting opioid to have effect before, durin, aducmd(f)3(or)3(6-1( )]TJ 0[95( )Tj [

- **Stimulation** causes the generalized excitement seen in horses and cats (often together with tachycardia); euphoria (tends to abuse); dysphoria (unwanted), and vomiting and defecation (unwanted).
- All opioids exert similar mode of action, but their activity varies depending on degree of affinity of the specific subtypes
- Mechanism of action
    - Mainly through its agonist activity at opioid receptors that results in modulation of

    - e.g., morphine, fentanyl, meperidine
- Antagonists
    - Drugs that competitively reverse (antagonize) the effects of agonists by preventing their access to a receptor.
    - They have low (or no) intrinsic activity at the receptors themselves
    - e.g., naloxone, nalmefene, naltrexone
- Partial agonists
    - Drugs that produce a submaximal response at a particular receptor type, even at high doses
    - The slope of dose-response curve is **less steep** that that of pure (full) agonist
    -

- Use with care in the horse (maximum 0. 1mg/ kg, preferably with sedation). Vomiting and defecation is a common side effect.
- It can be used at all perioperative stages.
- Inexpensive
- Time of onset after IV injection is 10 minutes; IM takes longer.
- Rapid IV injection causes more predictable histamine release (give it slow in this route)
- Doses of 0.1-0.3 mg/kg provide 4 h ours analgesia in the dog.
- Vomiting and defecation is not uncommon
- Urinary retention and constipation with prolonged usage

Oxymorphone

- It is μ agonist, and very similar to morphine; approximately 10 times more potent
- Vomiting is rare and produces some mild sedation
- us5J 0.001 Tadd3Tq4sjo0k37 0 Td( )Tj EMC /LBody <</MCID 38 >>BDC /C2_0 1 Tf 9.96 310

Methadone

- This is equipotent to morphine but gives more prolonged analgesia
- It causes less vomiting than morphine
- IV administration does not cause histamine release, so given this route for faster onset of drug action
- In horses, use at 0.1 mg/kg IM (IV can get them excited if not with sedatives); dogs and cats up to 0.5 mg/kg (but beware of respiratory depression)

Fentanyl

- μ agonist
- Very potent short acting quick onset analgesic
- Popular as intra-operative (and post-operative) CRI analgesic
- Transdermal patch offers advantage of prolonged duration of effect
- The patches come in several doses each accordingly recommended for different patient sizes
- They need to be applied at least 12 hours before the expected time of expected start of the operation to be effective in dogs (much faster onset in horses)
- However, once the plasma level is established the analgesia lasts for a few days.
- It does not cause histamine release
- Analgesia is dose related, but is accompanied by respiratory depression, and at high doses (up to 20 mcg/kg) its use may be only with O_2 and IPPV.
- In dogs, an infusion could start with a 3-5 mcg/kg as bolus followed by an infusion of 3 to 6 mcg/kg/hr. In cats, half of these doses can be used.
- Cardiovascular effects are minimal, but bradycardia may be seen
- It tends to cause excitement (manifested by walking) in the horse
- Other congeners (e.g., sufentanyl, remifentanyl, alfentanyl etc.) are too expensive for current veterinary use
- Carfentanyl, another analogue, is very very potent, and has been primarily used for darting wild lifes.
- Pre-mixed formulations (typically with butyrophenones; e.g., droperidol or fluanisone) may be available

Opioid antagonists

Naloxone

- It is pure antagonist
- It attaches to opioid receptors but have no effect
- The most common use is for reversing opioid induced sedation, respiratory depression, excitement.
- It is recommended agent for treating opioid overdose in man.
- Doses are 1 to 25 mcg/kg
- Naltrexone and nalmefene are longer acting pure antagonists, but have been used less than naloxone
- NB1. The duration of an antagonist may not be the same as that of the agonist, so the animal may return to sleep or respiratory depressant. (Close observation must be continued until the negative agonist effect fully wanes)

- NB2. If an opioid antagonist is used, it will reverse analgesia, and post-operative analgesia cannot be easily supplied with further opioids. The term **sequential analgesia** refers to the use of a partial agonist such as buprenorphine or nalbuphine to reverse a pure agonist such as morphine or fentanyl; the idea then being that some analgesia will remain. There are differing views as to whether this works in practice.

Opioid partial agonists

- Principles of using antagonists are similar to those relating to sedative antagonists
- It is important that the length of action of the agonists and antagonists are matched. Where opioid antagonists are used, analgesia is also reversed

Buprenorphine

- a μ partial agonist,

- It is 5 to 10 times more potent than morphine, but analgesic efficacy is much less (NB, do not confuse ‘potency’ with ‘efficacy’), and is not very effective in managing severe pain (exception is in bird as this seems provide a good analgesia).
- It provides mild sedation
- Cardiopulmonary effects are minimal
- With increasing dose there is a ceiling effect on respiratory depression (i.e., further increase do not increase respiratory depression)
- In dogs, 0.1 to 0.5 mg/kg, in cats, 0.1-0.4 mg/kg, and in horses, 0.02-0.05 mg/kg can be given IV, IM, or SQ. In the horse 0.1 mg/kg IV causes marked ‘walking behavior’. In dogs, IM is painful so beware of being bitten.
- It has a duration of action lasting approximately 2 hours

Pentazocine

- partial agonist, μ antagonist, has some activity at higher doses
- Frequently causes dysphoria, and is now only occasionally used
- It lasts approximately 1 to 2 hours
- Class IV drug

Use of opioids with sedatives

- When sedative and opio

- Carfentanyl and Etorphine are the most dangerous, but buprenorphine and fentanyl and its analogues are also effective by this route.
- Always must be handled by the veterinary professionals, and have emergency CPR kits and personnel who can perform the CPR available when handling the potent opioids.

Non steroidal anti-inflammatory drugs (NSAIDs).

- Until relatively recently it has been stated that NSAIDs were not effective in acute pain, and therefore (although useful later) were inadequate for immediate post-operative analgesia
- However, the modem potent NSAIDs such as carprofen and ketoprofen give useful post-operative analgesia.
- It is probable that they do not provide any surgical analgesia (carprofen does not reduce the MAC of halothane in dogs by an appreciable amount and they do not reduce the sensitivity to acute pain in analgesiometric tests in sheep).
- When compared with opioids for post-operative analgesia, the NSAIDs have the advantage of not decreasing consciousness or causing respiratory depression.
- Their anti-inflammatory action speeds healing and decreases pain.
- They are toxic, and tend to be cumulative.
- There are major species differences in elimination times and toxicity, and data and doses cannot be transposed between species.
- In general it is not advisable to administer these drugs intravenously during anesthesia, as the delay in elimination (through anesthetic induced cardiovascular depression) again increases toxicity.
- For all NSAIDs, never exceed the data sheet dose quantity or in frequency of administration.
- Flunixine meglumate (Banamin)
  - It has been used in the horse for some years.
  - There appear to be no problems with single dose IV use during anesthesia.
  - However, cumulation through multiple dosing can cause colitis, particularly in sick horses.
  - Overdose causes death. It has been shown to produce worth while analgesia during and after surgery when given as premedication.
  - Possible renal toxicity has been reported when it has been given to halothane anaesthetised dogs.
- Carprofen (Rimadyl)
  - It is very widely used analgesic in the dogs
  - Appears very effective at doses of 4mg/kg IV or SQ given at premedication or induction.
  - It is very long acting and has a much greater safety margin than has flunixine
  - The range available is enormous. For details of doses, half-lives, and toxicity (in dog and cat) see the physiology reference)
- Many NSAIDs may be given by the oral route, making them useful for the later stages of postoperative analgesia.
- Other useful perioperative NSAID analgesics include meloxicam, firocoxib, etodolac, and tepoxalin, and much of the toxicity is claimed to have been lessened due to their selective COX (particularly COX2) or LOX inhibitory action. For full details of the dosing and duration etc see the suggested references below

Alpha 2 adrenoceptor agonists

- It provides some effective analgesia, particularly effective in visceral pain in comparison to somatic pain
- When used during anesthesia it reduce the dose of other anesthetics required
- However, as analgesia is accompanied by sedation, in addition to all the other side effects their use for post-operative analgesia is not common.
- Alpha 2 agonists and opioid combination shows marked synergism, and it is possible to obtain all degrees of sedation and, in some cases, anesthesia with such combinations. No pre-prepared combinations exist.
- They can provide effective analgesia with minimal side effects when used by the epidural route.
- If antagonists have been used, analgesia is reversed alongside sedation.

NMDA antagonists

- Ketamine falls into this classification
- Tends to provide better somatic analgesia than visceral analgesia
- CRI infusion now commonly used in some severely painful animals, particularly in the ICU setting, but usually is combined with other analgesics and sedatives to enhance the analgesic efficacy and reduce dissociative side effects
- Amantadine has been used in man with varying success, and seems to have a place in controlling a neuropathic pain

Other adjunctive analgesic therapy

- Acupuncture
- Transcutaneous electrical stimulation (TENS)
- Physical therapy
- Antidepressant
- Anticonvulsants etc.
- For further information on these modalities see the references cited below

Conclusion

- What determines which agents and methods for pain control should be used would depend on the type of procedures, severity of pain and economic consideration for each individual circumstance.
- Our understanding of pain in its manifestation, mechanisms, assessment, and alleviation in animals is still, although improving, limited
- It is imperative to ensure continuous effort in establishing information on safe and appropriate use of analgesics in animals
- However, it must be remembered that no pharmacological intervention can supersede tender loving care which animals deserve most during their suffering